

CT Guided Protocol

For CT Scanner + Conventional Impressions

NO TEETH BEHIND CUSPIDS

1. Send Us The Digital RX Form

Completely fill out the online digital RX form found at www.nsequence.com or your case cannot be processed.

2. Patient CT Scan Wearing an Acrylic Duplicate Denture with Markers

Fabricated duplicate denture should have at least six markers. Scan the patient wearing the duplicate denture while biting on a bite registration at 0.3 voxel for 20 seconds (preferably 13cm field of view). Ensure the duplicate denture is fully seated in the patient's mouth. Reline the duplicate denture with Coe-Soft™ material to ensure a good seal. Alternatively, a denture teeth setup may be used as a better occlusal record.

3. Acrylic Duplicate Denture with Markers

Mail the duplicate denture in with the upper and lower VPS impressions.

4. Upper & Lower Impressions

Take full upper and lower VPS impressions of your patient. Be sure to capture vestibules and peripheral rolls, as anatomically distal and as accurately as possible.

5. Bite

Capture a full arch Blu-Mousse® bite of the patient relating upper and lower dentition with the duplicate denture in place at a closed position. If the patient's opposing is a denture, be sure to use the denture as the opposing.

6. Send Us Photos & Shades

Send us digital clinical photographs of the patient and the desired shade.

CT Guided Protocol

For CT Scanner + Intraoral Scanner

NO TEETH BEHIND CUSPIDS

1. Send Us The Digital RX Form

Completely fill out the online digital RX form found at www.nsequence.com or your case cannot be processed.

2. Patient CT Scan Wearing an Acrylic Duplicate Denture with Markers

Fabricated duplicate denture should have at least six markers. Scan the patient wearing the duplicate denture while biting on a bite registration at 0.3 voxel for 20 seconds (preferably 13cm field of view). Ensure the duplicate denture is fully seated in the patient's mouth. Reline the duplicate denture with Coe-Soft™ material to ensure a good seal. Alternatively, a denture teeth setup may be used as a better occlusal record.

3. Acrylic Duplicate Denture with Markers

Mail the duplicate denture in with the upper and lower VPS impressions.

4. Upper & Lower Digital Impressions

Use your COS scanner to take upper or lower digital impressions. Notify your service provider to send data to nSequence[®]. Be sure to capture vestibles and peripheral rolls as anatomically distal and accurately as possible. If not possible, VPS impressions are a suitable replacement.

5. Digital Bite Against Opposing

Use your intraoral scanner to take the bite of the denture against the opposing natural dentition.

6. Send Us Photos & Shades

Send us digital clinical photographs of the patient and the desired shade.